



A Novel likely Pathogenic No-Start Variant in HEY2 Associated with Electrical-Predominant Dilated Cardiomyopathy: A Case Report

Domenico Dell'Edera^{1,2*}, Allegra Donadio¹, Francesco La Rocca¹, Catia Fausto¹, Rosina Valicenti³

Abstract

Background: Inherited cardiomyopathies are genetically heterogeneous disorders and a major cause of heart failure, ventricular arrhythmias, and sudden cardiac death. The increasing use of next-generation sequencing has expanded the spectrum of disease-causing genes and improved genotype–phenotype correlations.

Case summary: We describe a 64-year-old Caucasian man with hypokinetic dilated cardiomyopathy (left ventricular ejection fraction 40–45%), frequent ventricular ectopy (approximately 18,000 premature ventricular complexes/24 h), diffuse left ventricular hypokinesia, and ascending aortic dilatation (47 mm). Genetic testing identified a previously unreported heterozygous no-start variant in HEY2 (NM_012259:c.-29_12del), involving the 5' untranslated region and the canonical translation initiation sequence. The variant is predicted to abolish normal translation initiation and was classified as likely pathogenic according to ACMG criteria (PVS1, PM2). To our knowledge, this variant has not previously been reported.

Discussion: This case expands the mutational spectrum of HEY2 and supports its role in inherited cardiomyopathy characterized by a predominant electrical phenotype with significant ventricular arrhythmias. Further functional studies and additional cases are needed to clarify the pathogenic mechanism and strengthen genotype–phenotype correlations.

Conclusion: This case expands the mutational spectrum of HEY2 by describing a previously unreported likely pathogenic no-start variant associated with dilated cardiomyopathy and a prominent ventricular arrhythmic phenotype. Although functional validation is lacking, the predicted disruption of translation initiation and the absence of the variant from population databases support its pathogenic role. This report further strengthens the evidence implicating HEY2 in inherited cardiomyopathies and highlights the value of comprehensive genetic testing for establishing genotype–phenotype correlations, improving risk stratification, and guiding family screening. Further studies are warranted to clarify the molecular mechanisms underlying HEY2-related cardiomyopathy.

Keywords: HEY2 gene; Upstream Open Reading Frames region (uORFs); 5' untranslated region (5'UTR region); Hypokinetic Dilated Cardiomyopathy; Cardiovascular Magnetic Resonance (CMR); Echocardiography.

Abbreviations

ACC: American College of Cardiology; ACM: arrhythmogenic cardiomyopathy. AHA: American Heart Association; AF: familial atrial

Affiliation:

¹Unit of Medical Genetics and Immunogenetics, “Madonna delle Grazie” Hospital, 75100 Matera, Italy

²Clinical Pathology and Microbiology Unit, “Papa Giovanni Paolo II” Hospital, 75025 Policoro (MT), Italy

³Preventive Cardiology Clinic, “Papa Giovanni Paolo II” Hospital, 75025 Policoro (MT), Italy

*Corresponding author:

Domenico Dell'Edera. Unit of Medical Genetics and Immunogenetics, “Madonna delle Grazie” Hospital, 75100 Matera, Italy.

Citation: Domenico Dell'Edera, Allegra Donadio, Francesco La Rocca, Catia Fausto, Rosina Valicenti. Cardiology and Cardiovascular Medicine. 10 (2026): 256-262.

Received: July 21, 2026

Accepted: July 28, 2026

Published: September 04, 2026

fibrillation; BrS: Brugada syndrome; CMP: cardiomyopathies; CPVT: catecholaminergic polymorphic ventricular tachycardia; DCM: dilated cardiomyopathy; ESC: European Society of Cardiology. HCM: hypertrophic cardiomyopathy; LQTS: long QT syndrome; PCCD: progressive familial cardiac conduction defect; RCM: restrictive cardiomyopathy; SQTS: short QT syndrome.

Introduction

Genetic cardiomyopathies are disorders of the heart muscle caused by mutations in genes that regulate the structure and function of the heart. Genetic arrhythmias are heart rhythm disorders caused by mutations in genes that alter the transmission of electrical impulses in the heart [1]. The classification of cardiomyopathies and genetic arrhythmias is fundamental for the diagnosis, treatment, and management of patients. The guidelines provided by the American College of Cardiology (ACC), the European Society of Cardiology (ESC), and the American Heart Association (AHA) provide distinct criteria, but generally agree on many of the categories [2,3]. According to the American College of Cardiology (ACC), cardiomyopathies are divided into four main categories: Dilated cardiomyopathy (DCM), Hypertrophic cardiomyopathy (HCM), Restrictive cardiomyopathy (RCM), Arrhythmogenic cardiomyopathy (ACM).

The American Heart Association (AHA) defines cardiomyopathies (CMP) as a “heterogeneous group of myocardial disorders characterized by mechanical and/or electrical dysfunction, which generally, but not always, manifest as inappropriate ventricular hypertrophy or dilation, resulting from various causes, frequently genetic.” The European Society of Cardiology defines cardiomyopathies as “a myocardial disorder characterized by structural and functional abnormalities of the heart muscle, in the absence of coronary artery disease, hypertension, valvular disease, and congenital heart disease sufficient to explain the observed myocardial abnormality.” As the most effective method for the diagnosis and daily management of patients, this classification system is more clinically oriented and categorizes heart muscle disorders based on ventricular morphology and function [4,5]. Based on the above, cardiomyopathies are classified as primary when the disease primarily affects the heart muscle, or as secondary when the disease is part of a broader disorder (syndromic pattern). Finally, primary cardiomyopathies can be classified as acquired, mixed (genetic and non-genetic), and hereditary. Cardiac arrhythmias are classified by the American College of Cardiology (ACC) as: Brugada syndrome (BrS), long QT syndrome (LQTS), short QT syndrome (SQTS), familial atrial fibrillation (AF), catecholaminergic polymorphic ventricular tachycardia (CPVT), and progressive familial cardiac conduction defect (PCCD). Hereditary cardiac arrhythmias

are rare conditions and often cause sudden death in young adults. They result from a series of mutations affecting various genes that control the function of ion channels and the proteins involved in their regulation. Since the alterations affect ion channels, they are also known as channelopathies [6-8]. A better understanding of genetics is important for identifying patients in the preclinical stage, who are unaware that they have a genetic defect that will cause them to develop cardiomyopathy or arrhythmia. This article reports on a study conducted on a 64-year-old man with hypokinetic dilated cardiomyopathy (EF 40-45%), massive ventricular/supraventricular arrhythmias (18,000 ventricular ectopic beats - VEB/24h), aortic dilatation (47 mm), diffuse hypokinesia of the left ventricle, and cardiac rhythm instability. The genetic study found for the first time the presence of the “no-start” variant NM_012259: c.-29_12del p.(?) in a heterozygous state in the Hairy Enhancer-of-split related with YRPW motif 2 (HEY2) gene located on the long arm of chromosome 6 (6q22.31) (dbSNP: rs759494126). According to the classification of the American College of Medical Genetics and Genomics (ACMG), this variant is classified as probably pathogenic (class IV, PVS1, PM2). In silico predictor values are: CADD 23.6, SpliceAI 0.00, Pangolin 0.0200, phyloP 3.97. The HEY2 protein acts as a transcriptional repressor and helps maintain ventricular electrophysiological identity, cell junction organization, and myocardial homeostasis. Therefore, its reduction can produce a cardiac phenotype that initially leads to electrophysiological rather than structural alterations.

Materials and Methods

Patient Information, History of Present Illness and Physical Examination

The subject is a 64-year-old man of Italian origin, normotensive (BP: 120/70 bpm), euglycemic, euthyroid, with mild dyslipidemia and no gastroesophageal reflux disease (GERD). The patient presented with recurrent palpitations, reduced exercise tolerance, and episodes of nocturnal palpitations. The patient had an echocardiogram before the CMR. The findings showed left ventricular dilation, moderate reduction in systolic function, and aortic root at the upper limits. These symptoms prompted Holter monitoring. The finding of a very high arrhythmic load justified subsequent CMR, performed to evaluate possible arrhythmic cardiomyopathy or fibrosis. The Holter dynamic electrocardiogram detected: a heart rate between 120 and 33 beats per minute (bpm), approximately 760 non-periodic premature ventricular contractions (PVCs), and approximately 9,000 ventricular extrasystoles occurring episodically in pairs. The extrasystoles were monomorphic, with a morphology consistent with a lower ventricular origin (left bundle branch block-like pattern). Cardiovascular

Magnetic Resonance (CMR) was performed using Siemens Magnetom Aerea 1.5 Tesla equipment, using Gadovist as the contrast agent (CA), administered intravenously at a dose of 0.15 mmol/kg [9]. Sequences were used to assess cardiac function (cine-SSFP), edema (TIRM, T2, and T2 mapping), fibrosis (T1 and T2 mapping), perfusion, and myocardial necrosis (PSIR and LGE) after administering the contrast agent. CMR revealed dilation of the aortic root (maximum diameter of approximately 47 mm) and ascending aorta (maximum diameter of approximately 41 mm), while the atria and heart valves were normal.

Diffuse segmental hypokinesia was observed with moderate reduction in global systolic function (EF=40%); no clear signal alterations suggestive of myocardial edema were detected in the TIRM T2 sequences; no alterations in myocardial relaxation times were observed in the T1 and T2 mapping sequences; no areas of scarring (fibrosis) or myocardial necrosis were observed in the PSIR and LGE sequences after contrast medium administration.

Clinical Findings and Investigations

These investigations led to the clinical conclusion that the subject had dilated cardiomyopathy, in the absence of clear alterations in tissue characterization. A genetic study was conducted to better understand the phenotype-genotype association. The molecular test we use is based on the principle of massive parallel sequencing (Next Generation Sequencing, NGS). This has made it possible, using a commercial kit (SOPHiA EXTENDED CARDIO SOLUTION™ Kit), to simultaneously study 128 genes involved in the hereditary transmission of certain cardiomyopathies and cardiac arrhythmias (the tested gene list is shown in Table 1).

Genetic Testing

Sample collection and genomic DNA extraction

Venous blood samples were collected in tubes containing ethylenediaminetetraacetic acid-K3 (EDTA-K3) as an anticoagulant after obtaining informed consent. Genomic DNA was isolated from leukocytes using the MagCore automated extractor (RBC Bioscience, Taiwan). The automated protocol ensures consistent yields and high purity,

minimizing operator variability. Extractions were performed according to the manufacturer's instructions, and purified genomic DNA was stored at -20°C until use. DNA purity and concentration were assessed by spectrophotometry using the Implen NanoPhotometer N60 (Implen GmbH, Germany), recording the absorbance ratios of 260/280 and 260/230 to check for protein contaminants. Quantitative DNA assay was performed using the Qubit 4.0 fluorimeter (ThermoFisher Scientific Holdings Europe Ltd, UK) with the Qubit dsDNA High Sensitivity kit, which allows for accurate measurement of double-stranded DNA content.

Library preparation and target enrichment

Target enrichment was performed using the Sophia Genetics Extended Cardio Solution (ExtCas) panel. This panel includes the simultaneous study of 128 genes related to the manifestation of genetically based cardiomyopathies and/or arrhythmias (Sophia Genetics, 401 Park Drive, 5TH Floor, Boston, MA 02215, USA). Library preparation was conducted according to the manufacturer's instructions, with minor operational modifications introduced to improve uniformity of coverage. Polymerase Chain Reaction (PCR) steps were performed on a VeritiPro Thermal Cycler (ThermoFisher Scientific Holdings Europe Ltd, UK). After amplification, PCR products were purified by magnetic separation to remove adapter dimers and small fragments. Library concentration was verified by fluorimetry, and fragment size distribution was analyzed using the Agilent 4200 TapeStation System (Agilent Technologies, 5301 Stevens Creek Blvd., Santa Clara, CA 95051, USA) to confirm the expected amplification profile before sequencing.

Sequencing and bioinformatics analysis

Normalized libraries were pooled in equimolar quantities and sequenced on the Illumina MiSeq Dx platform (Illumina Inc., USA), using the MiSeq Reagent Kit V2 in a 250-bp paired-end configuration. This setup ensured an optimal balance between sequencing depth and accuracy, allowing for the reliable identification of single nucleotide variants (SNVs) and small insertions or deletions. Each sequencing run included negative extraction controls and positive reference samples to monitor for contamination and evaluate

ABCC9, ACTA1, ACTC1, ACTN2, AKAP9, ALPK3, ANK2, ANKRD1, APOA1, ATP2A2, BAG3, CACNA2D1, CACNAIC, CACNB2, CALM1, CALM2, CALM3, CALR3, CASQ2, CAV3, CHRM2, CRAYB, CSRP3, CTF1, CTNNA3, DES, DMD, DOLK, DPP6, DSC2, DSG2, DSP, DTNA, EMD, EYA4, FGF12, FHL1, FHL2, FKTN, FLNC, GAA, GATA4, GATA6, GATADI, GJA1, GJA5, GJC1, GLA, GPD1L, HCN4, HEY2, HFE, JPH2, JUP, KCNA5, KCNAB2, KCND3, KCNE1, KCNE2, KCNE3, KCNE5, KCNH2, KCNJ2, KCNJ5, KCNJ8, KCNQ1, LAMA4, LAMP2, LDB3, LMNA, MYBPC3, MYH6, MYH7, MYL2, MYL3, MYLK2, MYOM1, MYOZ2, MYPN, NEBL, NEXN, NKX2-5, NOS1AP, NPPA, NUP155, PDLIM3, PKP2, PLN, PRDM16, PRKAG2, PSEN1, PSEN2, PTPN11, RAF1, RANGRF, RBM20, RYR2, SCN10A, SCN1B, SCN2B, SCN3B, SCN4B, SCN5A, SCO2, SGCD, SLC8A1, SLMAP, SNTA1, STRN, SURF1, TAZ, TBX20, TBX5, TCAP, TGFβ3, TMEM43, TMPO, TNNC1, TNNI3, TNNT2, TPM1, TRDN, TRPM4, TRPM7, TTN, TTR, VCL.

Table 1: The tested gene list.

overall run performance. FASTQ files were automatically generated using MiSeq Reporter software. Primary data processing, including demultiplexing, adapter trimming, and alignment to the GRCh37 (hg19) human reference genome, was performed using Sophia DDM software V7 (Sophia Genetics, Switzerland). Variant calling and annotation were performed using Sophia Genetics' proprietary algorithms, integrating curated databases such as ClinVar, COSMIC, and dbSNP. Quality control parameters (mean depth of coverage, uniformity, and percentage of on-target reads) were analyzed for each sample to ensure data reliability and reproducibility.

Discussion

Molecular testing identified the presence of the heterozygous variant c.-29_12del p.(?) in the Hairy Enhancer-of-split related with YRPW motif 2 (HEY2, NM_012259) gene. The Sophia DDM software V7 (Sophia Genetics, Switzerland), using the criteria of the American College of Medical Genetics and Genomics (ACMG), classifies this variant as “probably pathogenic” (class IV, PVS1, PM2) (Figure 1).

The deletion observed includes the loss of 31 nucleotides falling within the 5' untranslated region (5'UTR) and 10 nucleotides falling within the Upstream Open Reading Frames (uORF) region (nt: ATGAAGCGCC) (Figure 2) [10].

This gene encodes a member of the basic helix-loop-

helix (bHLH) transcription factor family, hairy and enhancer of split-related (HESR). The encoded protein is localized in the nucleus and interacts with a histone deacetylase complex to repress transcription. The expression of this gene is induced by the Notch signal transduction pathway [11]. Data in the literature show that reduction Hey2 expression in adult patients with cardiomyopathy (CM) causes contractile dysfunction and mortality [12], as well as in patients with dilated cardiomyopathy (DCM). Increased HEY2 expression has also been reported in patients with heart failure [13]. RNA-seq confirmed that there is a reduced expression of the mutated allele resulting from the heterozygous deletion in the HEY2 gene (RNAwt: 80%, RNA_{del}: 20%). The most striking finding in this case study is not the 40% EF, but approximately 760 nonperiodic ventricular extrasystoles (PVCs) and approximately 9,000 ventricular extrasystoles that occurred episodically in pairs. This suggests widespread electrical instability. Experimental models of reduced HEY2 function have observed altered expression of sodium channels, potassium channels, and proteins involved in impulse propagation. In the long run this leads to the development of PVC-induced cardiomyopathy (chronic extrasystole, functional remodeling, reduction of ejection fraction). When observed it is partly confirmed by a positive family history: the proband (family tree III2) shows that the paternal grandmother (family tree I2) had an cardioverter defibrillator fitted (ICD) (documentation not available as she died at the age of seventy-five); the father of the




Figure 1: HEY2 gene (NM_012259), variant c.-29_12del p.(?) in heterozygous. The “Sophia DDM software”, using the criteria of the American College of Medical Genetics and Genomics (A.C.M.G.), classifies this variant as “probably pathogenic” (class IV, PVS1, PM2).



Figure 2: The deletion involves the loss of 31 nucleotides in the 5'UTR region and 10 nucleotides in the upstream open reading frame (NT: ATGAAGCGCC). The deletion removes a regulatory sequence that consequently leads to increased expression of the HEY2 protein (IGV software).




Figure 3: Pedigree. Proband (II2) identified by the arrow. Square: male; circle: female; shaded: affected by cardiomyopathy and with a mutation in the HEY2 gene.

proband (family tree III1) died of suspected sudden cardiac death at the age of sixty-seven (SCD); the two sisters of the proband (family tree III1 and III3) underwent cardiological examinations and genetic testing, which found no cardiac abnormalities or the presence of the c.-29_12del p.(?) variant in the HEY2 gene (Figure 3).

Conclusion

In this study, we report for the first time the heterozygous variant NM_012259:c.-29_12del p.(?) in the Hairy Enhancer-of-split related with YRPW motif2 (HEY2) gene located on the long arm of chromosome 6 (6q22.31) and its association with “arrhythmogenic electrical-predominant cardiomyopathy associated with partial loss of HEY2 function”. The deletion eliminates the start codon and reduces gene activity; reduced HEY2 activity alters ventricular electrophysiological regulation; compare this with a high rate of ventricular extrasystoles; chronic arrhythmia induces moderate systolic dysfunction without evident fibrosis; the alteration of the Notch/HEY2 pathway itself may contribute to the observed aortic dilation. Table 2 provides a summary of the evidence provided by the ACMG.

Reduction of HEY2 activity alters the transmural gradient of ion channels; reduces sodium current density; and promotes afterdepolarizations. These mechanisms are perfectly compatible with a phenotype of frequent ventricular extrasystoles, predominant electrical arrhythmias, and systolic dysfunction secondary to tachyarrhythmia. HEY2 is part of the Notch pathway, which is also involved in vascular morphogenesis. Although no direct data exist on aortic dilation, the hypothesis is biologically plausible as an inference based

on the role of the Notch pathway in vascular wall stability. The presence of a ventricular arrhythmic phenotype in our proband is consistent with the genetic association between HEY2 and Brugada; the modulation of SCN5A by HEY2; and the reduction of sodium current in Hey2+/- models. The results reported here suggest a likely pathogenic effect of this variant and could serve as a springboard for functional studies aimed at better understanding the molecular pathways involved.

Table 2: Summary of ACMG evidence.

Parameter	Detail
HEY2 gene	<i>Hairy/Enhancer-of-split related with YRPW motif 2</i>
Genomic location	Chromosome 6q22.31
Variant type	No-start deletion: <i>NM_012259: c.-29_12del p.(?)</i>
Affected region	5'UTR (-31 nt) + uORF (-10 nt)
Molecular effect	Loss of start codon → absence of translation → <i>loss-of-function</i>
ACMG classification	Class IV – Likely Pathogenic (PVS1 + PM2 + PP3 + PP4)
Population frequency	Absent from gnomAD, ClinVar, HGMD
Biological role	Cardiac development, ventricular differentiation, electrical conduction
Structural phenotype	Hypokinetic DCM (EF 40–45%), diffuse LV hypokinesia
Electrical phenotype	18,000 PVCs/24h, complex ventricular and supraventricular arrhythmias
Aortic findings	Aortic dilation 47 mm
Genotype–phenotype correlation	Consistent with HEY2 loss-of-function: cardiomyopathy, arrhythmias, aortic abnormalities
Literature evidence	HEY2 linked to structural defects, arrhythmias, CHD, aortic anomalies
Clinical implications	Arrhythmic risk, DCM progression, family screening, tailored follow-up

Funding: No funding was received for this work.

Availability of data and materials: Not applicable.

Ethics approval and consent to participate: This study complies with the ethical standards of the institutional and/or national research committee and with the 1964 Declaration of Helsinki and its later amendments. Written informed consent was obtained from the patient for participation.

Consent for publication: Written informed consent was obtained from the patient and the patient's parent to write

and publish this case report. A copy of the written consent is available for review by the Editor-in Chief of this journal.

Competing interests: The authors declare no competing interests.

Conflicts of interest: The authors declare that they have no conflict of interest.

Acknowledgments

We thank the Volunteer Organizations (ODV) Gian Franco Lupo and Anima Mundi. The authors would like to thank their families for their help and participation in the study.

Author Contributions

Conceptualization: DD, AD; Methodology: DD; Investigation FL; CF: Writing original draft preparation; DD, and RV; Figure preparation RV and AD; Review and editing, DD, and RV; Supervision DD.

References

- Hayesmoore JB, Bhuiyan ZA, Coviello DA, et al. EMQN: Recommendations for genetic testing in inherited cardiomyopathies and arrhythmias. *Eur J Hum Genet* 31 (2023): 1003-1009.
- Maximilian Buja L, Jagdish Butany. *Cardiovascular Pathology* (2022).
- Hershberger RE, Givertz MM, Ho CY, et al. Genetic Evaluation of Cardiomyopathy-A Heart Failure Society of America Practice Guideline. *J Card Fail* 24 (2018): 281-302.
- Chou C, Chin MT. Genetic and Molecular Mechanisms of Hypertrophic Cardiomyopathy. *Int J Mol Sci* 24 (2023): 2522.
- Elliott P, Schunkert H, Bondue A, et al. Integration of genetic testing into diagnostic pathways for cardiomyopathies: a clinical consensus statement by the ESC Council on Cardiovascular Genomics. *Eur Heart J* 46 (2025): 344-353.
- Corrado D, Anastakis A, Basso C, et al. Proposed diagnostic criteria for arrhythmogenic cardiomyopathy: European Task Force consensus report. *Int J Cardiol* 395 (2024): 131447.
- Fu DG. Cardiac Arrhythmias: Diagnosis, Symptoms, and Treatments. *Cell Biochem Biophys* 73 (2015): 291-296.
- Hauer RN, Aliot E, Block M, et al. Working Group on Arrhythmias and Working Group on Cardiac Pacing. Indications for implantable cardioverter defibrillator (ICD) therapy. Study Group on Guidelines on ICDs of the Working Group on Arrhythmias and the Working Group

- on Cardiac Pacing of the European Society of Cardiology. Eur Heart J 22 (2001): 1074-1081.
9. Pugliese L, Luciano A, Chiochi M. The Role of Cardiac Magnetic Resonance Imaging in the Management of Hypertrophic Cardiomyopathy. J Cardiovasc Dev Dis 12 (2025): 189.
 10. Robinson JT, Thorvaldsdottir H, Turner D, Mesirov JP. igv.js: an embeddable JavaScript implementation of the Integrative Genomics Viewer (IGV). Bioinformatics 39 (2023): btac830.
 11. She P, Gao B, Li D, et al. The transcriptional repressor HEY2 regulates mitochondrial oxidative respiration to maintain cardiac homeostasis. Nat Commun 16 (2025): 232.
 12. van Walree ES, Dombrowsky G, Jansen IE, et al. Germline variants in HEY2 functional domains lead to congenital heart defects and thoracic aortic aneurysms. Genet Med 23 (2021): 103-110. Epub 2020 Aug 21. Erratum in: Genet Med 24 (2022): 965.
 13. Das S, Frisk C, Eriksson MJ, et al. Transcriptomics of cardiac biopsies reveals differences in patients with or without diagnostic parameters for heart failure with preserved ejection fraction. Sci Rep 9 (2019): 3179.



This article is an open access article distributed under the terms and conditions of the [Creative Commons Attribution \(CC-BY\) license 4.0](https://creativecommons.org/licenses/by/4.0/)